

Information and Documentation:

Data Elements within Rethink's Medical Necessity Assessment for ABA and Autism



Contents

Introduction to Rationale Behind Use of AI within theMedical Necessity Assessment	.3
Overview of Data Elements Collected	.5
References Supporting the Validity of the Fields Within the MNA	.6
References Validating the Use of Unsupervised Machine Learning (ML) to Identify Patient Clusters in Individuals with Autism Using Variables Collect within the MNA	11
References on Modeling Patient Progress as a Function of Hours of ABA and other Variables:	13
RethinkFirst Publications:	14



Introduction to Rationale Behind Use of AI within the Medical Necessity Assessment

Predictability is a common concern in the delivery of healthcare services. Patients typically want to know how likely it is that each of several treatment options will be effective (e.g., Meyer et al., 2021); and they want to know the potential treatment length and cost relative to their personal budget and calendar so they can plan and budget accordingly (e.g., Bhargava & Loewenstein, 2015; Han et al., 2012). Likewise, insurance companies often want to know how potential treatment length and cost so they can forecast expenses and plan strategically for the future (e.g., Papanicolas et al., 2018). Finally, healthcare practitioners want to know potential treatment length, the possible range in which outcomes can improve, and the resources it will require so they can manage things such as caseloads, waitlist communication, hiring, and business operations (e.g., WHO, 2015).

It is through this lens that the current work sits relative to one area of healthcare in the delivery of Applied Behavior Analysis (ABA) services for individuals with autism spectrum disorders (ASD). The prevalence of ASD diagnosis has increased in the previous 20 years from 1 in 150 to 1 in 44 children (Centers for Disease Control and Prevention, 2021). As a result, stakeholders (patients, parents, payors, and providers) are increasingly aware of the variability in cost, treatment duration, and patient outcomes. For example, the average annual cost of treatment is estimated to range anywhere between \$17,227 to \$130,182 (Autism Speaks, 2022; Eldevik et al., 2009); and individuals diagnosed with autism and their families might receive ABA for between 5 to 40 hours per week for a duration ranging between 18 months to five years (Larsson, 2012). Such variability in cost and duration of treatment following an ASD diagnosis adds significant uncertainty to all involved (De Groot & Thurik, 2015; Han et al., 2012; Meyer et al., 2021).

Compounding this uncertainty around increasing amounts of ABA service delivery is a lack of standardization for ABA dosage as a function of patient presentation at intake and throughout the duration of services. At the time of writing, the process of determining optimal dosage of ABA for each patient is seemingly subjective. Past research suggests a patient's age, symptom severity, historical treatment duration, personal/family needs, current abilities, and the overall goals of therapy should play a role in how many hours per week of ABA an individual might need (e.g., APBA, 2020; Granpeesheh et al., 2009). However, no known quantitative methodology with robust empirical support allows clinicians to translate those variables into a precise recommendation of the optimal number hours of ABA. Thus, a tool that provides reliable and precise predictions based on patient data and published best-practice evidence would help decrease the subjectivity and uncertainty around ABA treatment.

In addition to decreased subjectivity and reduced uncertainty around ABA treatment, board certified behavior analysts (BCBAs) arguably have an ethical obligation to improve in this area. The Behavior Analyst Certification Board (BACB) Ethics Code for Behavior Analysts includes several guidelines that speak directly to this topic (BACB, 2020). For example, Guideline 2.08 requires discussing the scope of treatment with a client before starting services; Guideline 3.01 requires BCBAs to identify and act upon opportunities that lead to avoidable harm or wasteful allocation of resources; and, Guideline 3.12



requires BCBAs to advocate for appropriate services. Here, leveraging a reliable, precise, and evidencebased method to recommend ABA dosage allows the BCBA to accurately communicate the scope of likely treatment; why they believe benefits will be maximized while avoiding wasteful allocation of resources; and to know that the amount and level of behavioral services they are recommending is databased.

Past researchers have sought to improve predictability and aid BCBAs in their ethical obligations by quantitatively modeling the relationship between the hours that patients contact ABA and the resulting outcomes. For example, Linstead and colleagues (2017a) modeled the relationship between treatment intensity and mastered learning objectives for 726 children aged 1.5-12 years who received community-based behavioral intervention services. Linear regression and neural network models led to r2 values of .35 and .60, respectively. Similarly, in a separate study by Linstead et al. (2017b), they used linear regression to model the relationship between treatment intensity/duration and goals mastered within specific skill domains for 1,468 individuals with ASD. Across goal domains, r2 values ranged between .50 (social goals) and .67 (motor goals). As a final example, Ostrovsky and colleagues (2022) used a series of t-tests, effect size measures, and Pearson correlations to quantify the relationship between changes in standardized assessment scores as a function of hours of ABA and modality of supervision. For the 178 individuals included in the study, they observed that clinically significant improvements in function were independent of the hours of ABA received.

Though broader limitations to this (and other) past studies are described in more detail below, the variance accounted for by these models was likely impacted by several factors. First, the overall sample sizes in these studies were relatively small2. Second, as noted by the authors of several studies, models have often been built without accounting for the heterogeneity that comprises the broad spectrum that is ASD and that was unlikely to be fully captured in the studied samples. A final limitation to past work in this realm (noted by Ostrovsky et al., 2022) is the assumption of a linear relationship between hours per week of ABA and progress made. Dose-response curves in other areas of behavioral healthcare are often decidedly nonlinear (e.g., Dews, 1955; Levy et al., 2020; Zoladz & Diamond, 2009). Thus, modeling approaches that more flexibly account for nonlinear relationships may perform better (e.g., the neural network outperforming linear regression in Linstead et al., 2017a).



Overview of Data Elements Collected

At a broad level, RethinkFirst collects data relevant to business and clinical operations related to applied behavior analysis (ABA) therapy. The resulting extensive database involves hundreds of tables and tens of thousands of unique data fields. Currently, the RethinkFirst database contains data from 322,000+ patients with autism spectrum disorders and related developmental disabilities; from 97,000,000+ hours of ABA therapy; and containing 700,000,000 data points on patient progress from ABA therapy.

The Medical Necessity Assessment currently captures the following clinical information:

- Patient demographics (e.g., age, gender, race, socioeconomic status),
- Caregiver demographics (e.g., socioeconomic status, highest education),
- Patient disease/disorder related information (e.g., diagnostic assessment results, criterion-referenced assessment results),
- Idiosyncratic symptom presentation,
- Specific behaviors targeted for change via ABA therapy and their corresponding developmental domain (e.g., social interaction, communication, restricted/repetitive behavior),
- Comorbid medical and behavioral conditions/concerns,
- Social determinants of health (SDOH),
- Barriers to treatment progress (e.g., illness, cancellations, waitlists),
- History of contact with ABA therapy,
- History of contact with supplemental therapies (e.g., speech-language pathology, occupational therapy, psychological services),
- Patient outcomes (e.g., goals mastered, change in assessment scores)

Operationally, these data include information such as:

- Amount of ABA-related hours requested (e.g., 97151-97158; 0362T, 0373T),
- Amount of supplemental therapy hours requested (e.g., 97533, 92507),
- Rate and timing of skills and behavior assessments,
- Structure and content of individual ABA therapy sessions (i.e., greater detail on ABA "dose" than hours contacted),
- Treatment decision-variability among providers (e.g., goal definitions and pairing, time in treatment),
- Coordination of care,
- Staff-to-patient ratio / caseload management,
- Continuity of care (e.g., staff turnover, rate of staffing change per patient),
- Waitlist durations / time to treatment, Rate of treatment reduction / time in treatment,
- Customer selected quality measures (e.g., structure, process, outcome measures)
- Individual-level and organization-level provider performance comparisons/benchmarking,



• Individual-level and organizational-level performance variability

References Supporting the Validity of the Fields Within the Medical Necessity Assessment

- Agency for Healthcare Research and Quality (2014). Therapies for Children with Autism Spectrum Disorder: Behavioral Interventions Updates. Executive Summary. Behavior Analyst Certification Board (2014) Applied behavior analysis treatment of autism spectrum disorder: Practice guidelines for healthcare funders and managers (2nd ed.).
- Anderson, Angelika & Moore, Dennis & Rausa, Vanessa & Finkelstein, Simon & Pearl, Shaun & Stevenson, Mitchell. (2017). A Systematic Review of Interventions for Adults with Autism Spectrum Disorder to Promote Employment. Review Journal of Autism and Developmental Disorders. 4. 10.1007/s40489-016-0094-9.
- Anderson DK, Maye MP, Lord C. Changes in maladaptive behaviors from mid-childhood to young adulthood in autism spectrum disorder. Am J Intellect Dev Disabil. 2011 Sep;116(5):381-97. doi: 10.1352/1944-7558-116.5.381. PMID: 21905806; PMCID: PMC3185304.
- Behavior Analyst Certification Board (2022) Definition of Applied Behavior Analysis. Retrieved from https://www.bacb.com/wp-content/uploads/2020/05/BACB-Fact-Sheet
- Blacklock, K. & Perry, A. (2012). The Effectiveness of Intensive Behavioural Intervention In Children with Autism Over the Age of 6 Years: A Matched-Sample Controlled Study.
- Bronsard, Guillaume & Botbol, Michel & Tordjman, Sylvie. (2010). Aggression in Low Functioning Children and Adolescents with Autistic Disorder. PloS one. 5. e14358. 10.1371/journal.pone.0014358.
- Brown, H. D. (2007). Principles of language learning and teaching. White Plains, NY. Pearson Longman.
- Brugha TS, Doos L, Tempier A, Einfield S, & Howlin, P. Outcome measures in intervention trials for adults with autism spectrum disorders: A systematic review of assessments of core autism features and associated emotional and behavioral problems. International Journal of Methods in Psychiatric Research. 2015; 24(2):99-115.
- Center for Disease Control (2016) Autism and Developmental Disabilities Monitoring Network Community Report on Autism. Retrieved from
 - http://www.cdc.gov/ncbddd/autism/documents/comm-report-autism-full-report.pdf
- Centers for Medicare & Medicaid Services (2022). Glossary for Medically Necessary. Retrieved from https://www.cms.gov/apps/glossary/search.asp?Term=medically+necessary&Language=English &SubmitTermSrch=Search
- Centers for Medicare & Medicaid Services (2019). Medicare Program Integrity Manual. Reasonable and Necessary Provisions in Local Care Determinations. CMS 100-08, ch. 13, § 13.5.4. Retrieved from <u>https://www.cms.gov/Regulations-and-</u>

Guidance/Guidance/Manuals/Downloads/pim83c13.pdf

Council on Children with Disabilities, American Academy of Pediatrics Council on Children with



Disabilities, & Council on Children with Disabilities. (2007). Role of the medical home in family-centered early intervention services. Pediatrics, 120(5), 1153-1158.

- Council on Children with Disabilities, Section on Developmental Behavioral Pediatrics, Bright Futures Steering Committee, Medical Home Initiatives for Children with Special Needs Project Advisory Committee, Section on Developmental Behavioral Pediatrics, Medical Home Initiatives for Children with Special Needs Project Advisory Committee, et al. (2006). Identifying infants and young children with developmental disorders in the GUIDANCE FOR UNDERSTANDING TREATMENT DOSAGE 19 medical home: An algorithm for developmental surveillance and screening. Pediatrics, 118(1), 405-420.
- Davis-Wilson, D., (2018) Establishing Clinician Guidance for Understanding Treatment Dosage
 Recommendations for ABA Therapy and their Role in Utilization Management and Medical
 Necessity Determination: A Review of the Literature. Aspen Behavior Consulting. For use by
 Rethink Autism, Inc. Pg 1-21.
- Deliens, G., Leproult, R., Schmitz, R., Destrebecqz, A., & Peigneux, P. (2015). Sleep disturbances in autism spectrum disorders. Review Journal of Autism and Developmental Disorders, 2(4), 343-356.
- Denboba, D., McPherson, M. G., Kenney, M. K., Strickland, B., & Newacheck, P. W. (2006). Achieving family and provider partnerships for children with special health care needs. Pediatrics, 118(4), 1607-1615.
- Department of Health & Human Services: Centers for Medicare & Medicaid Services. (2014) CMCS Informational Bulletin: Clarification to Medicaid Coverage Services to Children with Autism; July 7, 2014.
- Department of Health & Human Services: Centers for Medicare & Medicaid Services. (2014) Medicaid and CHIP FAQs: Services to Address Autism; September 24, 2014.
- Diagnostic and Statics Manual of Mental Disorder, 5th Edition. (2013) American Psychiatric Association: Autism Spectrum Disorders, pg 50-59.
- Dupuis, Annie & Moon, Michael Jongho & Brian, Jessica & Georgiades, Stelios & Levy, Tomer & Anagnostou, Evdokia & Nicolson, Rob & Schachar, Russell & Crosbie, Jennifer. (2021). Concurrent Validity of the ABAS-II Questionnaire with the Vineland II Interview for Adaptive Behavior in a Pediatric ASD Sample: High Correspondence Despite Systematically Lower Scores. Journal of Autism and Developmental Disorders. 51. 10.1007/s10803-020-04597-y.
- Eikeseth, S., (2008). Early and Intensive Behavioral Intervention: Addressing Behavior Excesses and Deficits in Young Children with ASD. Autism Advocate 53(1), 15-19.
- Eldevik, S., Eikenseth, S., Jahr, E., & Smith, T. (2006). Effects of low-intensity behavioral treatment for children with autism and mental retardation. Journal of Autism and Developmental Disorders, 36(2):211 – 224.
- Eldevik, S., Hastings, R. P., Hughes, J. C., Jahr, E., Eikeseth, S., & Cross, S. (2010). Using participant data to extend the evidence base for intensive behavioral interventions for children with autism.
 American Journal on Intellectual and Developmental Disabilities, 115(5), 381-405).
- Florean IS, Dobrean A, Păsărelu CR, Georgescu RD, Milea I. The Efficacy of Internet-Based Parenting



Programs for Children and Adolescents with Behavior Problems: A Meta-Analysis of Randomized Clinical Trials. Clin Child Fam Psychol Rev. 2020 Dec;23(4):510-528. doi: 10.1007/s10567-020-00326-0. Epub 2020 Sep 8. PMID: 32897527.

- Frechter, Y., Demirsoy, I., Cameron, M. J., & Wirtjes, P. (2022). Toward a Value-Based Care Model for Children with Autism Spectrum Disorder. *Open Access Journal of Behavioral Science & Psychology*, 5(1).
- Gabrielsen, T. P., Farley, M., Speer, L., Villalobos, M., Baker, C. N., & Miller, J. (2015). Identifying autism in a brief observation. Pediatrics, 135(2), e330-e338.
- Gerhardt, P.F., Lainer, I. Addressing the Needs of Adolescents and Adults with Autism: A Crisis on the Horizon. J Contemp Psychother **41**, 37–45 (2011). <u>https://doi.org/10.1007/s10879-010-9160-2</u>
- Granpeesheh, D., Dixon, D. R., Tarbox, J., Kaplan, A. M., & Wilke, A. E. (2009). The effects of age and treatment intensity on behavioral intervention outcomes for children with autism spectrum disorders. Research in Autism Spectrum Disorders. Retrieved from: https://thestaracademy.co.za/wp-content/uploads/2016/03/Granpeesheh.pdf
- Granpeesheh, D., Tarbox, J., & Dixon, D. (2009). Applied behavior analytic interventions for children with autism: A description and review of treatment research. Annals of Clinical Psychiatry, 21(3): 162-173.
- Goldman, S. E., Richdale, A. L., Clemons, T., & Malow, B. A. (2012). Parental sleep concerns in autism spectrum disorders: Variations from childhood to adolescence. Journal of Autism and Developmental Disorders, 42(4), 531-8
- Golnik, A., Maccabee-Ryaboy, N., Scal, P., Wey, A., & Gaillard, P. (2012). Shared decision making:
 Improving care for children with autism. Intellectual and Developmental Disabilities, 50(4), 322-331.
- Golnik, A., Scal, P., Wey, A., & Gaillard, P. (2012). Autism-specific primary care medical home intervention. Journal of Autism and Developmental Disorders, 42(6). GUIDANCE FOR UNDERSTANDING TREATMENT DOSAGE 20
- Green, G., Brennan, L. C., & Fein, D., (2002). Intensive Behavioral Treatment for a Toddler at High Risk for Autism. Behavior Modification, 26(1): 69-102.
- Harris, Sandra & Handleman, Jan. (2000). Age and IQ at Intake as Predictors of Placement for Young Children with Autism: A Four- to Six-Year Follow-Up. Journal of autism and developmental disorders. 30. 137-42. 10.1023/A:1005459606120.
- Hendricks, Dawn & Wehman, Paul. (2009). Transition From School to Adulthood for Youth With Autism Spectrum Disorders. Focus on Autism and Other Developmental Disabilities. 24. 77-88.
 10.1177/1088357608329827.
- Hill, A., Zuckerman, K., & Fombonne. E. (2015). Obesity and Autism. Pediatrics. 2015; 136; 1051. DOI: 10.1542/peds.2015-1437.
- Hoffman, C. D., Sweeney, D. P., Lopez-Wagner, M., Hodge, D., Nam, C. Y., & Botts, B. H. (2008). Children with autism: Sleep problems and mothers' stress. Focus on Autism and Other Developmental Disabilities, 23(3), 155-165.
- Hyman SL, Levy SE, Myers SM; Council on Children with Disabilities, Section on Developmental and



Behavioral Pediatrics. Identification, Evaluation, and Management of Children With Autism Spectrum Disorder. Pediatrics. 2020 Jan;145(1):e20193447. doi: 10.1542/peds.2019-3447. Epub 2019 Dec 16. PMID: 31843864.

- Kenny, Lorcan & Cribb, Serena & Pellicano, Elizabeth. (2018). Childhood Executive Function Predicts Later Autistic Features and Adaptive Behavior in Young Autistic People: a 12-Year Prospective Study. Journal of Abnormal Child Psychology. 47. 10.1007/s10802-018-0493-8.
- Larsson, E. V. (2012). Applied Behavior Analysis (ABA) for Autism: What is the Effective Age Range for Treatment? The Lovaas Institute for Early Intervention. Retrieved from Larsson 2021 Is ABA and EIBI an Effective Treatment for Autism - History of Independent Panels (behavior.org)
- Liptak, G. S., Stuart T., & Auinger P. (2006). Health care utilization and expenditures for children with autism: data from U.S. national samples. Journal of Autism and Developmental Disorders, 36(7), 871-879.
- Linstead E, Dixon DR, Hong E, Burns CO, French R, Novack MN, Granpeesheh D. An evaluation of the effects of intensity and duration on outcomes across treatment domains for children with autism spectrum disorder. Transl Psychiatry. 2017 Sep 19;7(9):e1234. doi: 10.1038/tp.2017.207. PMID: 28925999; PMCID: PMC5639250.
- Lokhandwala, Tasneem, Khanna, Rahul (2012). Hospitalization Burden Among Individual with Autism. Journal of Autism and Developmental Disorders, 42:95-104
- Lord, C. & McGee, J. P., (2001). Educating Children with Autism. National Research Council. National Academy Press, Washington, D.C.
- Lovaas, O. I., (1987). Behavioral treatment and normal educational and intellectual functioning in young autistic children. Journal of Consulting and Clinical Psychology, 55(1): 3-9. Retrieved from https://doi.apa.org/doiLanding?doi=10.1037%2F0022-006X.55.1.3.
- Love, J. R., Carr, J. E., & Almason, S. M. (2009). Early and intensive behavioral intervention for autism: A survey of clinical practices. Research in Autism Spectrum Disorders, 3(2): 421 -428.
- Mandell, D., Xie, M., Knashawn H., Lawer, L., McCarthy, & M., Marcus S. (2012). The interplay of Outpatient Services in Psychiatric Hospitalizations Among Medicad-Enrolled Children with Autism Spectrum Disorders. Health and Human Services Public Access.
- Matson, J., Turygin, N., Beighley, J., Rieske, R.,, Tureck, K., & Matson, M. (2001). Applied Behavior Analysis in Autism Spectrum Disorders: Recent strengths, pitfalls. Research in Autism Spectrum Disorders.144-150
- Matsumura N, Fujino H, Yamamoto T, Tanida Y, Ishii A, Tatsumi A, Nakanishi M, Tachibana M, Mohri I, Okuno H. Effectiveness of a Parent Training Programme for Parents of Adolescents with Autism Spectrum Disorders: Aiming to Improve Daily Living Skills. Int J Environ Res Public Health. 2022 Feb 18;19(4):2363. doi: 10.3390/ijerph19042363. PMID: 35206550; PMCID: PMC8871773.
- Medical Comorbidities in Autism Spectrum Disorders. A Primer for Health Care Professionals and Policy Makers. (2013). Treating Autism Publications. Retrieved from https://nationalautismassociation.org/pdf/MedicalComorbiditiesinASD2013.pdf
- Myers, S.M., & Plauche Johnson, C. (2007) reaffirmed (2014). Management of Children with Autism Spectrum Disorders. Pediatrics, 120(5), 1162-1182.
- Nolan, K. W., Orlando, M., & Liptak, G. S. (2007). Care coordination services for children with special health care needs: Are we family-centered yet? Families, Systems, and Health, 25(3), 293-306.



- O'Rourke, C., Linden, M., & Bedell, G. (2020). Scoping Review of Interventions to Promote Social Participation in Adolescents and Young Adults with Neurodisability. *The Open Journal of Occupational Therapy*, 8(1), 1-22. <u>https://doi.org/10.15453/2168-6408.1647</u>
- Orinstein AJ, Helt M, Troyb E, Tyson KE, Barton ML, Eigsti IM, Naigles L, Fein DA. Intervention for Optimal outcome in children and adolescents with a history of autism. J Dev Behav Pediatr. 2014 May;35(4):247-56. doi: 10.1097/DBP.0000000000000037. PMID: 24799263; PCID:PMC4487510
- Peters-Sheffer, N., Didden, R., Korzilius, H., & Sturmey, P. (2011). A meta-analytic study on the effectiveness of comprehensive ABA-based programs for children with Autism Spectrum Disorders. Research in Autism Spectrum Disorders, 5(1): 60-69.
- Pellecchia, M., Connell, J. E., Beidas, R. S., Xie, M., Marcus, S. C., & Mandell, D. S. (2015). Dismantling the active ingredients of an intervention for children with autism. Journal of Autism and Developmental Disorders, 45(9), 2917- 2927.
- Powell PS, Pazol K, Wiggins LD, Daniels JL, Dichter GS, Bradley CB, Pretzel R, Kloetzer J, McKenzie C, Scott A, Robinson B, Sims AS, Kasten EP, Fallin MD, Levy SE, Dietz PM, Cogswell ME. Health Status and Health Care Use Among Adolescents Identified With and Without Autism in Early Childhood Four U.S. Sites, 2018-2020. MMWR Morb Mortal Wkly Rep. 2021 Apr 30;70(17):605-611. doi: 10.15585/mmwr.mm7017a1. PMID: 33914722; PMCID: PMC8084123.
- Ramey, C.T & Ramey, S.L. (1998). Early intervention and early experience. American Psychologist, February, 109-120. GUIDANCE FOR UNDERSTANDING TREATMENT DOSAGE 21.
- Remington, B., Hastings, R. P., Kovshoff, H., Espinosa, F. D., Jahr, E., Brown, T., Alsford, P., Lemaic, M., & Ward, N. (2007). Early intensive behavioral intervention: Outcomes for children with autism and their parents after two years. American Journal on Mental Retardation, 112(6): 418-438.
- Roth ME, Gillis JM, DiGennaro Reed FD. A meta-analysis of behavioral interventions for adolescents and adults with autism spectrum disorders. Journal of Behavioral Education 2014; 23(2): 258-286
- Sallows, G. O., & Graupner, T. D. (2005). Intensive behavioral treatment for children with autism: Four year outcome and predictors. American Association on Mental Retardation, 110(6), 417-438.
- Schickedanz A, Halfon N. Evolving Roles for Health Care in Supporting Healthy Child Development. Future Child. 2020 Fall;30(2):143-164. PMID: 33875912; PMCID: PMC8053141.
- Scott, M., & Condillac , R. (n.d.). Treatment Adherence in ABA: Practices and Opinions of Behaviour Analysts. *Research Special Interest Group Abstracts* , 23(2), 119–119.
- Scott M. Myers, Chris Plauché Johnson, the Council on Children With Disabilities; Management of Children With Autism Spectrum Disorders. *Pediatrics* November 2007; 120 (5): 1162–1182. 10.1542/peds.2007-2362
- Smith, S. W., Daunic, A. P., & Taylor, G. G. (2007). Treatment Fidelity in Applied Educational Research:
 Expanding the Adoption and Application of Measures to Ensure Evidence-Based
 Practice. Education and Treatment of Children, 30(4): 1-14.
- Stango, A. (2017). A Response to "Applied Behavior Analysis, autism, and occupational therapy: a search for understanding," as published in the American Journal of Occupational Therapy by Welch and Polatajko. Science in Autism Treatment, 14(1), 34-36.
- Tager-Flusberg H, Kasari C. Minimally verbal school-aged children with autism spectrum disorder: the neglected end of the spectrum. Autism Res. 2013 Dec;6(6):468-78. doi: 10.1002/aur.1329. Epub 2013 Oct 7. PMID: 24124067; PMCID: PMC3869868.



- Tiura M, Kim J, Detmers D, Baldi H. Predictors of longitudinal ABA treatment outcomes for children with autism: A growth curve analysis. Res Dev Disabil. 2017 Nov;70:185-197. doi: 10.1016/j.ridd.2017.09.008. Epub 2017 Sep
- Trasande, L. Chatterjee, S. (2009). The impact of obesity on health service utilization and costs in childhood. Obesity (Silver Spring). ;17(9):1749–1754.
- Virues-Ortega, J. (2010). Applied behavior analytic intervention for autism in early childhood: meta analysis, meta-regression and dose-response meta-analysis of multiple outcomes. Clinical Psychology Review, 30(4): 387-399.
- Vismara LA, Colombi C, Rogers SJ. Can one hour per week of therapy lead to lasting changes in young children with autism? Autism. 2009 Jan;13(1):93-115. doi: 10.1177/1362361307098516. PMID: 19176579; PMCID: PMC6992381.
- Volkmar, F., Siegel, M., Woodbury-Smith, M., King, B., McCracken, J., State, M., and the Academy of Child and Adolescent Psychiatry Committee on Quality Issues. (2014). Practice Parameter for the Assessment and Treatment of Children and Adolescents with Autism Spectrum Disorder. Journal of the American Academy of Child and Adolescent Psychiatry, 53(2) 237-257.
- Wasserman, Joan & Palmer, Richard & Gomez, Marcia & Berzon, Rick & Ibrahim, Said & Ayanian, John.
 (2019). Advancing Health Services Research to Eliminate Health Care Disparities. American Journal of Public Health. 109. S64-S69. 10.2105/AJPH.2018.304922.
- Wattanawongwan, Sanikan & Ganz, J & Pierson, Lauren & Hong, Ee Rea & Yllades, Valeria & Dunn, Claudia & Foster, Margaret. (2022). Quality Review of Social-Communication Interventions for Adolescents and Adults with ASD: Maintenance, Generalization, and Social Validity. Education and Training in Autism and Developmental Disabilitites. 57. 67-91.
- Woodman AC, Smith LE, Greenberg JS, Mailick MR. Contextual Factors Predict Patterns of Change in Functioning over 10 Years Among Adolescents and Adults with Autism Spectrum Disorders. J Autism Dev Disord. 2016 Jan;46(1):176-189. doi: 10.1007/s10803-015-2561-z. PMID: 26319253; PMCID: PMC4707127.

References Validating the Use of Unsupervised Machine Learning (ML) to Identify Patient Clusters in Individuals with Autism Using Variables Collect within Rethink's Medical Necessity Assessment

- Bitsika, V., Arnold, W. M., & Sharpley, C. F. (2018). Cluster analysis of autism spectrum disorder symptomatology: Qualitatively distinct subtypes or quantitative degrees of severity of a single disorder? *Research in Developmental Disabilities*, 76, 65-75.
- Broome, K., McCabe, P., Docking, K., Doble, M., & Carrigg, B. (2023). Speech Development Across Subgroups of Autistic Children: A Longitudinal Study. *Journal of Autism and Developmental Disorders*, 53(2570–2586). <u>https://doi.org/10.1007/s10803-022-05561-8</u>
- Christensen, D. L., Van Naarden Braun, K., Baio, J., et al. (2018). Prevalence and Characteristics of Autism



Spectrum Disorder Among Children Aged 8 Years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2012. MMWR / November 16, 2018 / Vol. 65 / No. 13. US Department of Health and Human Services/Centers for Disease Control and Prevention.

Dillon, E. F., Kanne, S., Landa, R. J., Annett, R., Bernier, R., Bradley, C., Carpenter, L., Kim, S. H., Parish Morris, J., Schultz, R., & Wodka, E. L. (2023). Sex Differences in Autism: Examining Intrinsic and Extrinsic Factors in Children and Adolescents Enrolled in a National ASD Cohort. Journal of Autism and Developmental Disorders, 53(1305–1318). <u>https://doi.org/10.1007/s10803-021-05385-y</u>

- Doshi-Velez, F., Ge, Y., & Kohane, I. (2014). Comorbidity Clusters in Autism Spectrum Disorders: An Electronic Health Record Time-Series Analysis. *Pediatrics*, 133 (1), e54-e63.
- Eaves, L. C., Ho, H. H., & Eaves, D. M. (1994). Subtypes of Autism by Cluster Analysis. *Journal of Autism and Developmental Disorders*, 24(1), 3-22.
- Edwards, D. J., Perlman, A., & Reed, P. (2012). Unsupervised Categorization in a Sample of Children with Autism Spectrum Disorders. *Research in Developmental Disabilities*, 33, 1264-1269.
- Gardner-Hoag, J., Novack, M., Parlett-Pelleriti, C., Stevens, E., Dixon, D., & Linstead, E. (2021). Unsupervised Machine Learning for Identifying Challenging Behavior Profiles to Explore Cluster-Based Treatment Efficacy in Children With Autism Spectrum Disorder: Retrospective Data Analysis Study. JMIR Medical Informatics, 9(6), e27793.
- Hart, C. M., & Curtin, S. (2023). Trajectories of Vocabulary Development in Children with Autism Spectrum Disorder Across Multiple Measures. *Journal of Autism and Developmental Disorders*, 53(1347–1362).
- Hotez, E., & Shea, L. (2023). Prioritizing Social Determinants of Health in Public Health Surveillance for Autism. *Pediatrics*, 151(2), e2022059541.
- Klopper, F., Testa, R., Pantelis, C., & Skafidas, E. (2017). A cluster analysis exploration of autism spectrum disorder subgroups in children without intellectual disability. *Research in Autism Spectrum Disorders*, 36, 66-77.
- Lingren T, Chen P, Bochenek J, DoshiVelez F, Manning-Courtney P, Bickel J, et al. (2016)

Electronic Health Record Based Algorithm to Identify Patients with Autism Spectrum Disorder. PLoS ONE 11(7): e0159621. doi:10.1371/journal.pone.0159621

- Matta, J., Zhao, J., Ercal, G., & Obafemi-Ajayi, T. (2018). Applications of node-based resilience graph theoretic framework to clustering autism spectrum disorders phenotypes. *Applied Network Science*, 3(38).
- Parlett-Pelleriti, C.M., Stevens, E., Dixon, D. *et al.* Applications of Unsupervised Machine Learning in Autism Spectrum Disorder Research: a Review. *Rev J Autism Dev Disord* (2022). <u>https://doi.org/10.1007/s40489-021-00299-y</u>
- Rescorla, L. (1988). Cluster analytic identification of autistic preschoolers. *Journal of Autism and Developmental Disorders*, 18(4), 475-492.
- Rivard, M., Mestari, Z., Morin, D. *et al.* Cluster Analysis of Clinical Features of Children Suspected to Have Neurodevelopmental Disorders. *J Autism Dev Disord* **53**, 2409–2420 (2023). https://doi.org/10.1007/s10803-022-05533-y

Rixon, L., Hastings, R. P., Kovshof, H., & Bailey, T. (2021). Sibling Adjustment and Sibling Relationships



Associated with Clusters of Needs in Children with Autism: A Novel Methodological Approach. *Journal of Autism and Developmental Disorders*, 51(4067–4076).

- Sevin, J. A., Matson, J. L., Coe, D., Love, S. R., Matese, M. J., & Benavidez, D. A. (1995). Empirically Derived Subtypes of Pervasive Developmental Disorders: A Cluster Analytic Study. *Journal of Autism* and Developmental Disorders, 25(6), 561-578.
- Stevens, E., Dixon, D. R., Novack, M. N., Granpeesheh, D., Smith, T., & Linstead, E. (2019). Identification and analysis of behavioral phenotypes in autism spectrum disorder via unsupervised machine learning. *International Journal of Medical Informatics*, 129, 29-36.
- Stevens, M. C., Fein, D. A., Dunn, M., Allen, D., Waterhouse, L. H., Feinstein, C., & Rapin, I. (2000).
 Subgroups of Children With Autism by Cluster Analysis: A Longitudinal Examination. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(3), 346-352.

References on Modeling Patient Progress as a Function of Hours of ABA and other Variables

- Eckes, T., Buhlmann, U., Holling, H.-D., & Möllmann, A. (2023). Comprehensive ABA-based interventions in the treatment of children with autism spectrum disorder – a meta-analysis. Psychiatry, 23(133).
- Eldevik, S., Hastings, R. P., Hughes, J. C., Jahr, E., Eikeseth, S., & Cross, S. (2009). Meta-Analysis of Early Intensive Behavioral Intervention for Children With Autism. Journal of Clinical Child & Adolescent Psychology, 38(3), 439-450.
- Fucà, E., Guerrera, S., Valeri, G., Casula, L., Novello, R. L., Menghini, D., & Vicari, S. (2023). Psychiatric Comorbidities in Children and Adolescents with High-Functioning Autism Spectrum Disorder: A Study on Prevalence, Distribution and Clinical Features in an Italian Sample. Journal of Clinical Medicine, 12, 677. <u>https://doi.org/10.3390/jcm12020677</u>.
- Kohli, M., Kar, A. K., Bangalore, A., & Prathosh, A. P. (2022). Machine learning-based ABA treatment recommendation and personalization for autism spectrum disorder: An exploratory study. Brain Informatics, 9(16).
- Linstead, E., Dixon, D. R., French, R., Granpeesheh, D., Adams, H., German, R., Powell, A., Stevens, E.,
 Tarbox, J., & Kornack, J. (2017). Intensity and Learning Outcomes in the Treatment of Children
 With Autism Spectrum Disorder. Behavior Modification, 41(2), 229–255.
- Linstead, E., Dixon, D. R., Hong, E., Burns, C. O., French, R., Novack, M. N., & Granpeesheh, D. (2017). An evaluation of the effects of intensity and duration on outcomes across treatment domains for children with autism spectrum disorder. Translational Psychiatry, 7, e1234. doi:10.1038/tp.2017.207.
- Maglione, M. A., Gans, D., Das, L., Timbie, J., Kasari, C., and For the Technical Expert Panel, HRSA Autism Intervention Research – Behavioral (AIR-B) Network. (2012). Nonmedical Interventions for Children With ASD: Recommended Guidelines and Further Research Needs. Pediatrics, 130(Supplement 2), S169-S178.
- Maharjan, J., Garikipati, A., Dinenno, F. A., Ciobanu, M., Barnes, G., Browning, E., DeCurzio, J., Mao, Q., & Das, R. (2023). Machine learning determination of applied behavioral analysis treatment plan



type. Brain Informatics, 10(7).

- Ostrovsky, A., Willa, M., Cho, T., Strandberg, M., Howard, S., & Davitian, C. (Year). Data-Driven, Client Centric Applied Behavior Analysis Treatment-Dose Optimization Improves Functional Outcomes. World Journal of Pediatrics. Published online 17 November 2022.
- Préfontaine, I., Morizot, J., Lanovaz, M. J., & Rivard, M. (2023). Changes in Autistic Symptoms and Adaptive Functioning of Children Receiving Early Behavioral Intervention in a Community Setting: A Latent Growth Curve Analysis. Journal of Autism and Developmental Disorders, 53(901–917).
- Schwichtenberg, A., & Poehlmann, J. (2007). Applied behaviour analysis: Does intervention intensity relate to family stressors and maternal well-being? Journal of Intellectual Disability Research, 51(8), 598-605. doi:10.1111/j.1365-2788.2006.00940.x
- Shannon, J., Salomon, C., Chettiath, T., Abbas, H., & Taraman, S. (Year). Autism Spectrum Disorder and the Promise of Artificial Intelligence. Journal of Child & Adolescent Behavior, 10(1), 1000428. ISSN: 2375-4494.
- Tiura, M., Kim, J., Detmers, D., & Baldi, H. (2017). Predictors of longitudinal ABA treatment outcomes for children with autism: A growth curve analysis. Research in Developmental Disabilities, 70, 185-197.
- Weiss, J. A., Cappadocia, M. C., MacMullin, J. A., Viecili, M., & Lunsky, Y. (2012). The impact of child

problem behaviors of children with ASD on parent mental health: The mediating role of acceptance and empowerment. Autism, 16(3), 261–274.

RethinkFirst Publications

Please click the purple "i" icon by the Medical Necessity header on any screen within the MNA and the BH Help Center.